POMPEJUS et al.

Serial No. 09/582,779

COPY OF ALL CLAIMS

- An orotidine-5'-phosphate decarboxylase gene having the sequence SEQ ID
 NO: 1 or its homologs which have at least 80% homology with the sequence
 SEQ ID NO: 1.
- An orotidine-5'-phosphate decarboxylase gene having the sequence SEQ ID NO: 1 or its homologs, wherein the gene or its homologs derive from Ashbya gossypii.
- 3. An amino-acid sequence encoded by a gene or its homologs as claimed in claim1.
- 4. An amino-acid sequence as claimed in claim 3, which comprises an enzymatically active protein.
- 5. A gene construct comprising an orotidine-5'-phosphate decarboxylase gene having the sequence SEQ ID No: 1 or its homologs as claimed in claim 1, where the gene or its homologs is functionally linked to one or more regulatory signals.
- 6. A gene construct as claimed in claim 5, whose gene expression is increased by the regulatory signals.
- 7. A vector comprising a gene construct as claimed in claim 5.
- 8. A microorganism comprising at least one gene construct as claimed in claim 5.
- 9. A process for producing uracil-auxotrophic microorganisms, which comprises modifying an orotidine-5'-phosphate decarboxylase gene having the sequence SEQ ID NO: 1 or its homologs as claimed in claim 1 in such a way that the

POMPEJUS et al.

Serial No. 09/582,779

protein encoded by the gene is inactive, and introducing this modified gene into the microorganisms and integrating said gene by homologous recombination into the genome of the organisms, and subsequently selecting these microorganisms for resistance to 5-fluoroorotic acid.

- 10. A process for inserting DNA into microorganisms, which comprises inserting a vector which comprises an intact orotidine-5'-phosphate decarboxylase gene having the sequence SEQ ID NO: 1 or its homologs as claimed in claim 1 together with at least one other nucleic acid sequence, into a microorganism which is deficient in orotidine-5'-phosphate decarboxylase nucleic acid sequence having the sequence SEQ ID NO: 1 or its homologs as claimed in claim 1 together with at least one other nucleic acid sequence, into a microorganism which is deficient in orotidine-5'-phosphate decarboxylase nucleic acid sequences, and cultivating this microorganism on or in a culture medium without uracil.
- 11. A process as claimed in claim 10, wherein a linear DNA is used as vector.
- 12. A process as claimed in claim 10, wherein an Ashbya gossypii strain is used as microorganism deficient in orotidine-5'-phosphate decarboxylase genes.
- 13. A process as claimed in claim 10, wherein at least one gene of riboflavin synthesis is inserted as additional gene into the microorganism.
- 14. A process for selecting cells, said process comprising the step of transforming cells with a gene sequence or its homologs as claimed in claim 1.

POMPEJUS et al.

Serial No. 09/582,779

15. The process as claimed in claim 14 for Ashbya gossypii.